# **Imported Wheat Distillers**













A very palatable, high energy and protein feed, rich in digestible fibre, low in starch and providing a good source of bypass protein.

### **Typical Analysis** (on a dry matter basis)

Dry matter (%)	Energy (MJ ME/kg DM)	Crude protein (%)	Oil (%)	NDF (%)	Starch (%)	Sugar (%)	DUP (%)
92.0	13.4	34.8	7.0	31.5	2.0	1.0	12.0

## What are you trying to achieve?

Need	Feature	Benefit		
Drive intake	Highly palatable feed.	Can stimulate intakes of less palatable feeds, increasing milk and meat production.		
Reduce feed costs	High quality protein and a good source of bypass protein.	Allows ratios of soya and low protein concentrates to be replaced whilst providing similar energy and protein levels (usually at lower cost).		
Improve rumen efficiency	Distillery products contain high levels of yeast fragments particularly in the solubles fraction.	Stimulates rumen activity, promoting fibre digestion and overall feed efficiency.		
Minimise risk of acidosis	High proportion of the energy as digestible fibre.	Allows energy intakes to be increased without increasing the risk of acidosis associated with high starch feeds.		
Feeding flexibility	Pellet durability.	Suitable for use in 'in parlour', automated and floor feeding systems. Can be transferred to feeders via auger systems.		

The predicted responses (benefits) assume that the specified nutrient, physical or structural dietary components are limiting livestock performance in the current ration.

#### **Complementary Concentrate Feeds**

• **High starch feeds** e.g. cereals, maize meals, and confectionary and bakery products.



0845 355 9935 www.kwfeeds.co.uk

- Low protein feeds e.g. cereals, soya hulls and sugar beet products.
- Rumen bypass proteins e.g. SoyPass, Prototec



#### Recommended daily feed rates (per head basis)

Imported Wheat Distillers can be fed via automated feeders, top dressed or floor fed, and used individually or as part of a blend or TMR.

Milking Cows	Up to 4 (typically 3)kg			
Dry Cows	Up to 2 kg			
Replacement Heifers	Up to 3 kg and up to 35% of the DMI			
Calves (to 12 weeks)	Up to 1.5 kg and up to 25% of the DMI			
Growing Cattle	Up to 2.5 kg and up to 40% of the DMI			
Finishing Cattle	Up to 5 kg and up to 40% of the DMI			
Suckler Cows	Up to 4 (typically 2)kg			
# Ewes and Rams	Up to 1 (typically 0.5)kg			
# Hoggets and Lambs	Up to 0.75kg and up to 50% of the DMI			
# /Unlike some feeds fro	m the which Industry co products from his ethanol production do not			

<sup># (</sup>Unlike some feeds from the whisky Industry, co-products from bio-ethanol production **do not** contain high levels of copper).

DMI = dry matter index

#### Availability, handling and storage

Imported Wheat Distillers are available as bulk tipped or blown loads. Like all dry feeds, they should be stored in a secure shed, bunker, bin or hopper and kept cool, dry and free from vermin. Imported Wheat Distillers' should be used within 2 months of delivery (due to a high residual oil content).

#### **Additional information**

#### Method of production

Imported Wheat Distillers are a product of the bio-ethanol industry. Following the fermentation of wheat and the distillation of ethanol, they are obtained from drying solid residues of fermented grains and adding evaporated syrups (solubles).

#### **Quality Assurance**

Imported Wheat Distillers are FEMAS assured (or a recognised equivalent) product. Imported Wheat Distillers are listed under number 1.12.11 in the EU Catalogue of Feed Materials.

#### **Legal Disclaimer**

Suggested feeding rates are produced as a guide only and many other factors may have an overriding effect on animal response; no performance guarantee can be given. Rations should be carefully balanced for energy and protein, contain sufficient forage to maintain rumen function and be fortified with an appropriate vitamin and mineral supplement. Animals must have constant access to clean water.



0845 355 9935 www.kwfeeds.co.uk



## **Imported Wheat Distillers**

## **Detailed Typical Analysis** (fresh basis other than where stated)

Dry matter	%	92.0	Calcium	g/kg	1.50
Oil A	%	5.50	Magnesium	g/kg	2.50
Oil B	%	6.50	Phosphorus	g/kg	0.90
Crude protein	%	32.0	Potassium	g/kg	1.10
Crude protein: DM	%	34.8	Salt	g/kg	3.50
Fibre	%	7.50	Sodium	g/kg	7.00
Ash	%	5.00	Copper	mg/kg	12.0
ME* – in vivo	MJ/kg DM	13.4	Manganese	mg/kg	65.0
NDF	%	29.0	Selenium	mg/kg	0.15
Starch	%	1.84	Zinc	mg/kg	85.0
Sugar	%	0.92	Saturates	% of oil	19.0
ERDP-FiM*	% @ 6%	19.5	Monounsaturates	% of oil	19.0
DUP-FiM*	% @ 6%	11.0	PUFAs	% of oil	62.0
DUP digestibility	%	82.0	Long chain PUFAs	% of oil	0.00
sDM		0.27	Lysine	% of CP	2.00
aDM		0.70	Methionine	% of CP	1.40
bDM		0.21	Cysteine	% of CP	1.75
cDM		0.11	Histidine	% of CP	2.25
sN		0.30	Threonine	% of CP	3.20
aN		0.74			
bN		0.18			
		0.17			
cDM sN aN		0.11 0.30 0.74 0.18	Histidine	% of CP	2.25

